

## Homework 9 Solution

Section 3.5 ~ 3.7.

3.5.8. Find all solutions of  $f(f(x)) = x$  for  $f(x) = -x^3/2$  and determine which are fixed points of  $f(x)$  and which are 2-cycles.

$$f(f(x)) = f\left(-\frac{x^3}{2}\right) = -\frac{\left(-\frac{x^3}{2}\right)^3}{2} = \frac{x^9}{16}$$

$$f(f(x)) = x \Rightarrow \frac{x^9}{16} - x = 0 \Rightarrow x^9 - 16x = 0$$

$$\Rightarrow x(x^8 - 16) = 0 \Rightarrow x(x^4 - 4)(x^4 + 4) = 0 \Rightarrow x(x^2 - 2)(x^2 + 2)(x^4 + 4) = 0$$

$$\Rightarrow x(x - \sqrt{2})(x + \sqrt{2})(x^2 + 2)(x^4 + 4) = 0 \Rightarrow x = 0, \sqrt{2}, -\sqrt{2}$$

$$f(0) = 0, f(\sqrt{2}) = -\sqrt{2}, f(-\sqrt{2}) = \sqrt{2}$$

Therefore 0 is a fixed point of  $f(x)$ , and  $\{\sqrt{2}, -\sqrt{2}\}$  is a 2-cycle.

3.5.10. Find all fixed points of  $f(x) = -x^2 + 2x + 2$  and then use them to help find all points of period 2.

$$f(x) = x \Rightarrow -x^2 + 2x + 2 = x \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow x = \frac{1 \pm \sqrt{9}}{2}$$

$$\begin{aligned} f(f(x)) &= f(-x^2 + 2x + 2) = -(-x^2 + 2x + 2)^2 + 2(-x^2 + 2x + 2) + 2 \\ &= -x^4 + 4x^3 - 2x^2 - 4x + 2 \end{aligned}$$

$$f(f(x)) = x \Rightarrow -x^4 + 4x^3 - 2x^2 - 4x + 2 = x \Rightarrow x^4 - 4x^3 + 2x^2 + 5x - 2 = 0$$

Then  $x^2 - x - 2$  is a factor of  $x^4 - 4x^3 + 2x^2 + 5x - 2$ , and

$$x^4 - 4x^3 + 2x^2 + 5x - 2 = (x^2 - x - 2)(x^2 - 3x + 1)$$

$$x^2 - 3x + 1 = 0 \Rightarrow x = \frac{3 \pm \sqrt{5}}{2}$$

One can check that  $f\left(\frac{3 + \sqrt{5}}{2}\right) = \frac{3 - \sqrt{5}}{2}$  and  $f\left(\frac{3 - \sqrt{5}}{2}\right) = \frac{3 + \sqrt{5}}{2}$ . Therefore

$$\left\{ \frac{3 + \sqrt{5}}{2}, \frac{3 - \sqrt{5}}{2} \right\}$$

is a 2-cycle.

3.5.18. Given that  $a = 1$  is one point of a 2-cycle of  $f(x) = 2 - 2^x$ , find the other point  $b$  and determine the stability of the cycle.

$$f(1) = 2 - 2^1 = 0, \quad f(0) = 2 - 2^0 = 1$$

So  $b = 1$ .

3.5.22. The equation  $P_{n+1} = \frac{1}{P_n} + \frac{P_n}{2} - 1$  is a price model.

(a) Find the two equilibrium points and determine their stability.

$$f(x) = \frac{1}{x} + \frac{x}{2} - 1$$

$$x = f(x) \Rightarrow x = \frac{1}{x} + \frac{x}{2} - 1 \Rightarrow \frac{x}{2} - \frac{1}{x} + 1 = 0 \Rightarrow x^2 + 2x - 2 = 0$$

$$\Rightarrow x = \frac{-2 \pm \sqrt{12}}{2} = -1 \pm \sqrt{3}$$

$$f'(x) = -\frac{1}{x^2} + \frac{1}{2}$$

$$f'(-1 + \sqrt{3}) = -\frac{1}{(-1 + \sqrt{3})^2} + \frac{1}{2} \approx -1.3660 < -1$$

So  $-1 + \sqrt{3}$  is an unstable equilibrium point.

$$|f'(-1 - \sqrt{3})| = \left| -\frac{1}{(-1 - \sqrt{3})^2} + \frac{1}{2} \right| \approx 0.3660 < 1$$

Therefore  $-1 - \sqrt{3}$  is a stable fixed point.

(b) Use the results from part (a) to help find the points of the 2-cycle, and then determine the stability of that cycle.

$$f(x) = \frac{1}{x} + \frac{x}{2} - 1 = \frac{x^2 - 2x + 2}{2x}$$

$$f(f(x)) = f\left(\frac{x^2 - 2x + 2}{2x}\right) = \frac{2x}{x^2 - 2x + 2} + \frac{x^2 - 2x + 2}{4x} - 1$$

$$= \frac{x^4 - 8x^3 + 24x^2 - 16x + 4}{4x^3 - 8x^2 + 8x}$$

$$f(f(x)) = x \Rightarrow \frac{x^4 - 8x^3 + 24x^2 - 16x + 4}{4x^3 - 8x^2 + 8x} = x$$

$$\Rightarrow x^4 - 8x^3 + 24x^2 - 16x + 4 = 4x^4 - 8x^3 + 8x^2 \Rightarrow 3x^4 - 16x^2 + 16x - 4 = 0$$

Note that  $x^2 + 2x - 2$  is a factor of  $3x^4 - 16x^2 + 16x - 4$ . Indeed,

$$3x^4 - 16x^2 + 16x - 4 = (x^2 + 2x - 2)(3x^2 - 6x + 2).$$

$$3x^2 - 6x + 2 = 0 \Rightarrow x = \frac{6 \pm \sqrt{12}}{6} = \frac{3 \pm \sqrt{3}}{3}$$

$$\left| f'\left(\frac{3 + \sqrt{3}}{3}\right) f'\left(\frac{3 - \sqrt{3}}{3}\right) \right| = 0.5 < 1$$

Therefore the 2-cycle

$$\left\{ \frac{3 + \sqrt{3}}{3}, \frac{3 - \sqrt{3}}{3} \right\}$$

is stable.

- 3.5.30. (a) Suppose  $f^4(p) = p$  for some point  $p$  and some function  $f(x)$ . What are the possible periods that  $p$  might have?

Because the period is the smallest  $m$  so that  $f^m(p) = p$ , the period of  $p$  is at most 4. But 3 is impossible, because if  $f^3(p) = p$ , then  $p = f^4(p) = f(f^3(p)) = f(p)$  and  $p$  is a fixed point (so the period is 1). Therefore the possible periods are 1, 2, and 4.

- (b) Suppose  $f^{12}(p) = p$  for some point  $p$  and some function  $f(x)$ . What are the possible periods that  $p$  might have?

By the same reason, the period of  $p$  is at most 12. We claim that the period of  $p$  must be a divisor of 12, so 1, 2, 3, 4, 6, and 12 are the only possible cases. Indeed, if the period  $m$  of  $p$  is less than 12, then  $f^m(p) = p$ . If we divide 12 by  $m$ , then  $12 = qm + r$  for some  $q \geq 1$  and the remainder  $m > r \geq 0$ . Now  $p = f^{12}(p) = f^r(f^{qm}(p)) = f^r(p)$ , so the period of  $p$  is at most  $r$  unless  $r = 0$ . The first case is impossible because the period  $m$  is larger than  $r$ . Therefore  $r = 0$  and  $m$  is a divisor of 12.

- 3.5.32. For  $f(x) = 1 - 4|x|$ ,  $p_1 = -11/65$  is a point of an  $m$ -cycle. Find all other points of that cycle, and determine its period and stability.

$$f\left(-\frac{11}{65}\right) = 1 - 4\left|-\frac{11}{65}\right| = \frac{21}{65}$$

$$f\left(\frac{21}{65}\right) = 1 - 4\left|\frac{21}{65}\right| = -\frac{19}{65}$$

$$f\left(-\frac{19}{65}\right) = 1 - 4\left|-\frac{19}{65}\right| = -\frac{11}{65}$$

Therefore  $p = -11/65$  has period 3. Note that

$$f(x) = \begin{cases} 1 - 4x, & x > 0 \\ 1 + 4x, & x < 0. \end{cases}$$

So

$$f'(x) = \begin{cases} -4, & x > 0 \\ 4, & x < 0 \end{cases}$$

Thus

$$\left|f'(-\frac{11}{65})f'(\frac{21}{65})f'(-\frac{19}{65})\right| = 64 > 1$$

and the 3-cycle is unstable.

3.6.2. Convert  $P_{n+1} = rP_n(1 - P_n^2/C^2)$  into a one-parameter family.

Let  $x_n = P_n/C$ . Then

$$Cx_{n+1} = rCx_n(1 - x_n^2) \Rightarrow x_{n+1} = rx_n(1 - x_n^2)$$

3.6.8. Find all positive fixed points of  $f_c(x) = \frac{cx^2}{x^2 + 1}$  and their intervals of existence for  $c > 0$ .

$$\begin{aligned} f_c(x) = x &\Rightarrow \frac{cx^2}{x^2 + 1} = x \Rightarrow cx^2 = x^3 + x \Rightarrow x^3 - cx^2 + x = 0 \\ &\Rightarrow x(x^2 - cx + 1) = 0 \\ &\Rightarrow x = 0, x = \frac{c \pm \sqrt{c^2 - 4}}{2} \end{aligned}$$

Because  $c > 0$ ,  $\frac{c + \sqrt{c^2 - 4}}{2} > 0$ . Furthermore,  $c^2 > c^2 - 4$  implies  $c > \sqrt{c^2 - 4}$ . So  $\frac{c - \sqrt{c^2 - 4}}{2} > 0$  as well. But these real solution exist only if  $c^2 - 4 \geq 0$  or equivalently,  $c \geq 2$  (note that  $c > 0$ ). Therefore the interval of existence is  $[2, \infty)$ .

3.6.12. Find the interval of stability of the fixed point 0 for  $r > 0$  of  $f_r(x) = rx^2(1 - x)$ .

$$f'_r(x) = 2rx - 3rx^2 \Rightarrow |f'_r(0)| = 0 < 1.$$

Therefore the fixed point 0 is always stable, so the interval of stability is  $(0, \infty)$ .

3.6.22. Find the 2-cycles and their intervals of existence and stability for  $a > 0$  of  $f_a(x) = -ax^3$ .

$$f_a(x) = x \Rightarrow -ax^3 = x \Rightarrow x(-ax^2 - 1) = 0 \Rightarrow x = 0$$

(Note that  $-ax^2 - 1 < 0$ .) So  $x = 0$  is the unique fixed point of  $f_a(x)$ .

$$f_a(f_a(x)) = f_a(-ax^3) = -a(-ax^3)^3 = a^4x^9$$

$$\begin{aligned} f_a(f_a(x)) = x &\Rightarrow a^4x^9 = x \Rightarrow x(a^4x^8 - 1) = 0 \Rightarrow x(ax^2 - 1)(ax^2 + 1)(a^2x^4 + 1) = 0 \\ &\Rightarrow x = 0, x = \frac{1}{\sqrt{a}}, x = -\frac{1}{\sqrt{a}} \end{aligned}$$

Since  $f_a(\frac{1}{\sqrt{a}}) = -\frac{1}{\sqrt{a}}$  and  $f_a(-\frac{1}{\sqrt{a}}) = \frac{1}{\sqrt{a}}$ ,  $\{\frac{1}{\sqrt{a}}, -\frac{1}{\sqrt{a}}\}$  is a 2-cycle. It exists for every  $a > 0$ . Therefore the interval of existence is  $(0, \infty)$ .

$$f'_a(x) = -3ax^2$$

$$|f'_a(\frac{1}{\sqrt{a}})f'_a(-\frac{1}{\sqrt{a}})| = 9 > 1$$

Therefore the 2-cycle is always unstable and the interval of stability is an empty-set.

3.6.28. For the threshold population model  $P_{n+1} = \frac{rP_n^2}{C}(1 - P_n/C)$ , when  $r$  is small the population will become extinct regardless of the initial population size  $P_0$ . When  $r$  is larger, although small initial populations still lead to extinction (because of the threshold), there is a positive equilibrium population.

(a) Find the smallest growth rate  $r$  for which a positive equilibrium population exists.

Set  $x_n = P_n/C$ . Then

$$Cx_{n+1} = rCx_n(1 - x_n) \Rightarrow x_{n+1} = rx_n(1 - x_n)$$

Let  $f_r(x) = rx(1 - x) = rx - rx^2$ .

$$f_r(x) = x \Rightarrow rx - rx^2 = x \Rightarrow rx^2 + (1 - r)x = 0 \Rightarrow x(rx + 1 - r) = 0$$

$$\Rightarrow x = 0, x = \frac{r - 1}{r}$$

Therefore the nonzero fixed point  $x = \frac{r-1}{r}$  is positive only if  $r > 1$ .

(b) Find the interval of  $r$ -values for which there exists a stable positive equilibrium population.

$$f'_r(x) = r - 2rx$$

$$f'_r(\frac{r-1}{r}) = r - 2r\frac{r-1}{r} = 2 - r$$

So  $|f'_r(\frac{r-1}{r})| < 1$  if and only if  $1 < r < 3$ . Therefore the interval of stability is  $(1, 3)$ .

3.7.2. Let  $f_r(x) = \frac{rx}{x + 5}$ .

(a) Find the interval of stability of the fixed point 0;

$$f'_r(x) = \frac{r(x + 5) - rx}{(x + 5)^2} = \frac{5r}{(x + 5)^2}$$

$$f'_r(0) = \frac{r}{5}$$

$$|f'_r(0)| = |\frac{r}{5}| < \Leftrightarrow r < 5$$

The interval of stability is  $(0, 5)$ .

- (b) Find the positive fixed point  $p(r)$  and its intervals of existence and stability;

$$f_r(x) = x \Rightarrow \frac{rx}{x+5} = x \Rightarrow rx = x(x+5)$$

$$\Rightarrow x^2 + 5x - rx = 0 \Rightarrow x(x+5-r) = 0 \Rightarrow x = 0, x = r-5$$

The nonzero fixed point  $p(r) = r-5$  is positive when  $r > 5$ . So the interval of existence is  $(5, \infty)$ .

$$f'_r(r-5) = \frac{5r}{(r-5+5)^2} = \frac{5}{r}$$

$$|f'_r(r-5)| = \frac{5}{r} < 1 \Leftrightarrow r > 5$$

Therefore the interval of stability is  $(5, \infty)$ .

- (c) Show that  $p(r)$  bifurcates from 0 at some parameter value  $r = r_0$ .

When  $r = 5$ ,  $p(5) = 5 - 5 = 0$ . At  $r = 5$ , the stable fixed point 0 becomes unstable and a new positive stable fixed point  $p(r)$  bifurcates from 0.

3.7.12. Let  $f_a(x) = 0.25 - ax^2$ .

- (a) Find the non-negative fixed point  $p(a)$  and its intervals of existence and stability;

$$f_a(x) = x \Rightarrow 0.25 - ax^2 = x \Rightarrow ax^2 + x - \frac{1}{4} = 0$$

$$\Rightarrow x = \frac{-1 \pm \sqrt{1+a}}{2a}$$

The positive solution is  $p(a) = \frac{-1 + \sqrt{1+a}}{2a}$  and it exists for every  $a > 0$ . Therefore the interval of existence is  $(0, \infty)$ .

$$f'_a(x) = -2ax \Rightarrow f'_a\left(\frac{-1 + \sqrt{1+a}}{2a}\right) = 1 - \sqrt{1+a}$$

So  $|f'_a(p(a))| = |1 - \sqrt{1+a}| < 1$  only if  $a < 3$ . Thus the interval of stability is  $(0, 3)$ .

- (b) Find the 2-cycle  $p_1(a), p_2(a)$  and its interval of existence and stability;

$$f_a(f_a(x)) = f_a(0.25 - ax^2) = 0.25 - a(0.25 - ax^2)^2 = \frac{1}{4} - a\left(\frac{1}{4} - ax^2\right)^2$$

$$= \frac{1}{4} - \frac{a}{16} + \frac{a^2}{2}x^2 - a^3x^4$$

$$f_a(f_a(x)) = x \Rightarrow a^3x^4 - \frac{a^2}{2}x^2 + x + \frac{a-4}{16} = 0$$

Note that  $ax^2 + x - \frac{1}{4}$  is a factor of  $a^3x^4 - \frac{a^2}{2}x^2 + x + \frac{a-4}{16}$ . Indeed,

$$a^3x^4 - \frac{a^2}{2}x^2 + x + \frac{a-4}{16} = (ax^2 + x - \frac{1}{4})(a^2x^2 - ax + \frac{4-a}{4}).$$

Now  $a^2x^2 - ax + \frac{4-a}{4} = 0$  has zeros

$$x = \frac{a \pm \sqrt{a^3 - 3a^2}}{2a^2} = \frac{1 \pm \sqrt{a-3}}{2a}.$$

$$p_1(a) = \frac{1 + \sqrt{a-3}}{2a}, \quad p_2(a) = \frac{1 - \sqrt{a-3}}{2a}$$

So they are two positive real roots when  $a > 3$  and  $1 - \sqrt{a-3} > 0$ , or equivalently,  $a < 4$ . Therefore the interval of existence (of a positive 2-cycle) is  $(3, 4)$ . Of course, the interval of existence (of a 2-cycle) is  $(3, \infty)$ .

$$|f'_a(\frac{1 + \sqrt{a-3}}{2a})f'_a(\frac{1 - \sqrt{a-3}}{2a})|$$

$$= |(1 + \sqrt{a-3})(1 - \sqrt{a-3})| = |1 - (a-3)| = |4 - a|$$

So it is less than one if  $3 < a < 5$ . Therefore the interval of stability (of a 2-cycle) is  $(3, 5)$ , and that of a positive 2-cycle is  $(3, 4)$ .

(c) Show that the 2-cycle bifurcates from  $p(a)$  at some parameter value  $a = a_0$ .

When  $a = 3$ ,  $p(3) = p_1(3) = p_2(3) = \frac{1}{6}$ . At  $a = 3$ , the stable fixed point  $p(3)$  becomes unstable and a stable positive 2-cycle  $p_1(a), p_2(a)$  bifurcates from  $p(a)$ .

3.7.16. Sketch the graph of  $p(a), p_1(a), p_2(a)$  from Exercise 12 together.

